

*SMAST Technical Report 21-03-06***Internal Tides in the Gulf of Maine, Part 1: Observations**


W. S. Brown, Z. Yu and R. Arena
 School for Marine Science and Technology
 University of Massachusetts Dartmouth
 New Bedford, Massachusetts
 (start date => 5 February 2004)

ABSTRACT

The Brown and Irish (1993) moored water property measurements in Georges Basin during 1986-87 showed strong semi-diurnal tidal variability. The Marsden (1986) moored current meter measurements on the north flank of Georges Bank indicated that the semi-diurnal internal tide was being generated there. The Georges Basin observations have been converted to inferred isopycnal displacement time series and analyzed in terms of the internal tidal content. The Garrett and Munk (1972) model - adapted for semi-diurnal frequencies - was used to estimate the velocity fields associated with the observed internal tidal elevations, that were very consistent with internal tidal generation on the north flank of Georges Bank. The Dartmouth finite element circulation model QUODDY (Lynch et al, 1996) was used to simulate the M2 internal tidal fields in the Gulf of Maine; simulations that show that standing the internal tidal waves are a prominent feature throughout the Gulf

BACKGROUND


The Gulf of Maine and the Bay of Fundy form a basin which is semi-enclosed by the relatively shallow Georges Bank. The Gulf of Maine system is connected to the North Atlantic Ocean mainly through the Northeast Channel and the Great South Channel (Figure 1). The Northeast Channel, with a sill depth of about 230 m, is flanked by the smaller Browns Bank to the northeast and the larger Georges Bank with a nominal depth of 60 m to the southwest. The Great South Channel with a sill depth of about 80 meters separates the southwestern end of Georges Bank from Cape Cod and Nantucket Shoals. There are three major basins separated by 200m sills within the gulf, namely the 280m deep Wilkinson and Jordan Basins and the 370m deep Georges Basin.



The geometry of the Gulf of Maine/Bay of Fundy system is the crucial factor in the amplified tidal response of the Gulf system. The characteristic or resonant period of the Gulf system is very near to the frequency of the semidiurnal tide (Garrett, 1972) which is forced by the tides in the North Atlantic. Thus the semidiurnal tidal response of the Gulf is amplified, while the diurnal tidal response is not. The M_2 semidiurnal harmonic constituent amplitude in the Gulf is four times greater than the next largest semidiurnal constituent, i.e., N_2 (Wright, 1986). The near-standing semidiurnal tidal wave response of the Gulf system ranges from about 40 cm along the seaward edge of Georges Bank to about 15 meters at the head of the Bay of Fundy.


The amplitudes and phases of tidal currents associated with the Gulf surface tide tend to be depth-independent (i.e. barotropic) except in the lowest part of the water column where friction is important. There is considerable lateral spatial structure of barotropic tidal currents because of the complex bathymetry of the Gulf (see Brown, 1984). Throughout much of the interior Gulf, the tidal currents are a relatively weak $O(10 \text{ cm/s})$. However, in regions of steep bathymetry, such as the north flank of Georges Bank, relatively stronger barotropic tidal currents $O(50 \text{ cm/s})$ are oriented across-isobath (Moody et al., 1984). Tidal currents in the Bay of Fundy are even stronger.

Internal or baroclinic tidal currents are generated by the interaction of the barotropic tidal currents and steep bathymetry in the presence of stratification. This interaction generates both a highly nonlinear internal solitary wave once every tidal cycle as well as a more regular internal tidal wave. The internal tidal wave appears to be generated continuously at depth along the north flank of Georges Bank and propagates toward the interior Gulf. Apparently, the internal solitary waves are generated as a lee wave associated with the incoming (i.e., flood) barotropic tidal current over the north flank of Georges Bank (Loder et al., 1992). When the flood tidal current begins to ebb, a solitary wave propagates toward the bank. As the solitary wave of isopycnal depression (or elevation) propagates away from a generation region, it can evolve into a packet of multiple depressions (Osborne and Burch, 1980). Internal wave packets also propagate away from Georges Bank (La Violette et al., 1990), as seen in the Space Shuttle photographs of the sea surface. These propagating internal wave packets are detected from space by the sea surface texture changes associated with the internal solitary wave surface flow convergence (ripples) and divergence (slicks) patterns (Sawyer, 1983; see Figure 2).



Marsden (1986) observed internal tides on the north flank of Georges Bank with moored current meters located at depths of 30, 40, and 75 meters in about 200 m of water (see Figure 3). He found that 98% of the current spectral energy density in the semidiurnal frequency band was contained in the mode in which the

across-isobath component was largest. Phase differences in the currents at adjacent levels indicated a downward energy propagation at the mooring site; consistent with the rays pattern emanating from the top of the slope. Horizontal wave length and phase speed were estimated to be 20 to 30 km and 0.40 to 0.70 m/s, respectively



As part of a Gulf-scale UNH study of the Gulf of Maine during 1986-87 (Brown and Irish, 1992, 1993) water property and circulation variability measurements--showing strong semidiurnal tidal influence--were made in Georges Basin. This paper describes those observations and their relation to the known internal tidal generation zone on the Bank. An analysis of the inferred isopycnal displacement time series will be presented. The Garrett and Munk (1972) model adapted for semidiurnal frequencies is used to estimate the internal tidal velocity fields associated with the inferred isopycnal displacement fields.

Observations

Brown and Irish (1992) describe the August 1968 through September 1987 Gulf of Maine study, during which a series of five "seasonal" hydrographic surveys were conducted. Water property and bottom pressure moorings were also deployed in Georges Basin, Wilkinson Basin, Jordan Basin, and the Northeast Channel, respectively.

This internal tides study focuses on the stratified water column time period between mid-August and mid-September 1986. Twenty mid-August CTD casts from Georges Basin (see [Figure 1](#)) were averaged to create a basin-averaged density profile ([Figure 4](#)). The robust stratification during this time period supported a strong internal tidal signal.

The 1986-87 GoM study Georges Basin moored observations were made at 42° 32.4'N, 67° 15.2'W ([Figure 2](#)). Temperature and conductivity sensor pairs on the Georges Basin mooring were located at 50m intervals between 20m, 70m, 120m, 170m, 220m and 270m depths. A bottom pressure/temperature instrument was located at 323m. The uninterrupted series of hourly measurements of temperature, salinity and bottom pressure (Brown and Irish, 1992) were obtained for the duration of the study. This analysis will focus on the Georges Basin mooring observations between 1800 UTC 11 August and 0100 UTC 18 September 1986.

Temperature and salinity measurements were used to compute a potential density anomaly time series. The time-averaged potential density anomalies at the depths of the moored sensor pairs in Figure 4 are very similar to those derived from a spatially-averaged hydrographic casts. Thus we were confident that the stratification structure did not vary significantly over the time period of interest.




Table 0. Summary statistics of the statistics, including maximum, minimum, mean and standard deviation of the UNH moored σ_θ at the levels indicated. The statistics of the constant σ_θ time series, which were generated for top and the bottom are also included.

name	max	min	mean	std dev
St0/ 0m sur	23	23	23	0
St1/ 20m	26.21	23.52	25.20	0.513
St2/ 70m	26.73	24.80	26.03	0.250
St3/ 120m	27.16	25.90	26.63	0.229
St4/ 270m	27.35	26.41	27.01	0.157
St5/ 220m	27.38	26.59	27.18	0.092
St6/ 270m	27.73	27.00	27.27	0.079
St6/ 323m bot	27.50	27.50	27.50	0

First, the moored density anomaly time series were high-pass filtered so as to severely attenuate the variability with periods longer than 36 hours. Then, a set of 17 selected isopycnal displacement time series were derived by way of a linear interpolation between the high-pass filtered density series (see Wallinga and Brown, 2004). The 10-day segment of a subset of the isopycnal displacement time series in Figure 5 clearly shows dominant semidiurnal frequency fluctuations, which were correlation at different depth throughout the water column. The statistics of the full suite of isopycnal displacements are presented in Table 1.

A partitioning of the energy density and variance preserving spectra of the isopycnal time series (Figure 6) show that about 76% of the isopycnal displacement energy at all depths is in the semidiurnal frequency band (0.0759 - 0.0848 cph; Table 2). Nonlinearities in the waveform are thought to be responsible for the lesser energy peaks of the higher harmonics. These results show that the semidiurnal frequency internal tide is dominant in Georges Basin. A harmonic analysis of the isopycnal displacement time series is consistent with these results.

The dominance of the correlated semidiurnal frequency variability of the water column is reflected in a time-domain empirical orthogonal function (T-EOF) analysis results in a first mode that explains about three fourths (74.8%) of the total variance. The mode-1 T-EOF structure has an amplitude maximum at 188 m depth with minimas at the surface and the bottom (Table 3); consistent with dynamical mode-1

characteristics. The mode-1 T-EOF amplitude time series (Figure 7) is clearly a semidiurnal signal. This mode is highly correlated (0.87) with the local barotropic pressure and lags it by about 0.8 hours.




Table 1. Summary of the basic statistics of selected isopycnals at the Georges Basin mooring site

Isopycnal Sigma-theta	Mean Depth (m)	Maximum Depth (m)	Minimum Depth (m)	Std. Dev. (m)
23.50	0.0	0.0	0.0	0.0
23.70	6.3	23.2	2.6	2.9
23.90	12.3	27.3	5.2	4.6
24.30	22.3	35.8	10.3	5.9
24.90	37.5	66.2	18.1	7.1
25.70	59.4	94.1	43.1	6.8
26.00	69.0	103.3	52.2	8.8
26.20	77.3	110.3	58.3	11.0
26.60	102.7	135.7	71.9	10.0
26.80	119.0	167.4	95.2	10.6
26.90	144.8	194.1	114.3	12.7
27.20	187.9	227.2	158.7	15.4
27.24	206.8	245.5	165.5	13.3
27.28	237.6	308.3	202.9	11.8
27.30	258.2	311.6	219.2	9.9
27.32	280.3	315.0	252.0	8.3
27.33	293.1	316.7	258.9	6.7
27.34	306.4	318.3	265.3	4.3
27.35	323.0	323.0	323.0	0.0





Table 2. A partition of the isopycnal displacement variance (σ^2) in Georges Basin. Note that the M_2 semidiurnal internal tide clearly dominates in the period band between 1 and 36 hours.

	σ^2 (m ²)	σ (m)	% Total σ^2
Total	1514.3	38.9	100.0
Mode-1 T-EOF	1132.7	33.7	74.5
SEMIDIURNAL BAND*	1049.3	32.4	69.3
M_2 TIDES	911.4	30.2	60.2
SEMIDIURNAL BAND	921.7	30.4	60.9
SD Mode-1 T-EOF	866.4	29.4	57.2



Analysis

A harmonic analysis of the isopycnal displacement series (Table 3) shows that about 60% of the total isopycnal displacement variance is explained by M₂ internal tidal response. The M₂ amplitudes have a maximum at 188 m and phases that are within +/- 15 degrees of a depth constant value of about 90 degrees G. The latter is the first-order structure of a classic dynamical mode-1 internal tide. Still, there is an approximate 30-degree phase increase from the surface to about 144 m depth. This phase structure indicated a slight downward internal wave phase velocity and an upward group or energy propagation velocity in this part of the water column.

The results of a frequency-domain EOF (F-EOF) analysis of the isopycnal displacement field are consistent with the above results. Perhaps it is not surprising that the mode-1 amplitude and phase structures (Figure 8; Table 3) are nearly identical to those of the M₂ harmonic amplitudes and phases. Mode-1 explains 94% of the variance in the semidiurnal frequency band of interest and 57% of the overall variance (Table 2).

Table 3. The structure of the isopycnal displacement series in terms of (a) the M₂ harmonic amplitude and Greenwich epoch; (b) the semidiurnal band mode-1 frequency-domain empirical orthogonal function (F-EOF) in terms of sinusoidal equivalent amplitude and Greenwich epoch phase; and (c) the model time-domain empirical orthogonal function (T-EOF)

Isopyc σ_t	Mean Depth (m)	Harmonic Amp (m)	Constants Phase (deg G)	Mode-1 F-EOF		T-EOF	
				Sin. Equ. Amp (m)	Phase (deg G)	Mode-1 Amp (m)	Mode-2 Amp (m)
-	0.0	-	-	-	-	-	-
23.70	6.3	2.79 +/-0.15	74 +/-3	2.56	74	2.25	
23.90	12.3	5.00 +/-0.17	75 +/-2	4.58	75	4.07	
24.30	22.3	7.08 +/-0.20	76 +/-1	6.51	76	5.91	
24.90	37.5	8.74 +/-0.30	82 +/-2	8.15	81	7.96	
25.70	59.4	8.58 +/-0.18	91 +/-1	8.13	89	8.53	
26.00	69.0	10.66 +/-0.21	94 +/-1	10.22	92	10.95	
26.20	77.3	13.84 +/-0.24	95 +/-1	13.26	93	14.03	
26.60	102.7	12.21 +/-0.29	99 +/-1	11.85	96	13.10	
26.80	119.0	11.76 +/-0.34	103 +/-1	11.77	101	13.68	
26.90	144.8	14.36 +/-0.37	104 +/-1	14.34	103	16.73	
27.20	187.9	17.23 +/-0.53	108 +/-2	17.06	107	20.36	
27.24	206.8	13.14 +/-0.72	107 +/-3	13.46	108	16.57	
27.28	237.6	11.37 +/-1.03	107 +/-5	11.07	110	13.87	
27.30	258.2	9.78 +/-0.98	107 +/-5	9.28	111	10.99	
27.32	280.3	7.62 +/-0.78	108 +/-5	7.17	112	8.21	
27.33	293.1	5.53 +/-0.65	107 +/-6	5.11	110	6.07	
27.34	306.4	2.83 +/-0.42	106 +/-8	2.50	109	3.21	
BP	323.0	0.82 +/-0.001	77 +/-0.1				




Figure 8. The first-mode empirical orthogonal function (EOF) in the [12 hour]⁻¹ frequency band. The isopycnal displacement equivalent sinusoidal amplitude and Greenwich epoch phases are presented. (a) The amplitude structure shows a general mid-depth maxima. (b) The phase structure is very nearly that of a classic mode-1 internal tide. Its depth-averaged phase is 100°G.

Analytical Model Internal Tides

The horizontal internal tidal currents in Georges Basin were modeled using the vertical displacement observations, the buoyancy frequency profile based on the basin-wide CTD observation (Figure 4) and the approximate depth-averaged phase from the harmonic analysis (101 degrees G). The model first-mode semidiurnal internal tide has a horizontal wavelength of 59.3 km and a phase speed of 132.5 1.0 cm/s. Given that the distance between the hypothesized generation region on Georges Bank north flank and the Georges Basin mooring is 38.7 km, it would take 8.08 hours (234 degrees phase lag) for a model internal tidal wave to reach the UNH mooring. This implies that the Georges Basin mooring is more than one half (0.65) model internal wavelengths from the assumed generation region.

The free parameter (C1) in the model formulation was set in accordance with the maximum amplitude of the observed vertical displacement. Figure 9 shows the model velocity and displacement amplitude structure based on Georges Basin parameters. The observed maximum amplitude of roughly 14.2 meters was located at a depth of 145 meters. The model amplitude maximum at a depth of 121 meters and was set to the observed maximum value of 14.2 m. The model M_2 internal tidal current hodographs in Figure 10 which is roughly (4 cm/s) twice as large as the along-isobath velocity. The kinematics of the model M_2 internal tide in Georges Basin (Figure 11) are those of a classic flat bottom ocean mode one internal gravity wave.

Figure 9. The first-mode M_2 internal tidal wave field in Georges Basin. (a) the horizontal longitudinal (solid) and transverse (dashed) velocity structure; (b) vertical velocity structure; and (c) vertical displacement structure.

Figure 10. Hodograph of the internal tidal horizontal velocities. Maximum surface velocities are 4 cm/s. The deeper water velocities taper off to order 1 cm/s.

Figure 11. Kinematics of a classic mode-1 M_2 internal tidal wave propagating from left (south) to right. Three isopycnals and the corresponding northward currents vectors for one tidal period. The convergence and divergence zones occur at the same time intervals. There is a 180-degree phase shift at the first e-folding depth scale of 121 m.

Analytical Model/Observations Comparison

The maximum in the observed isopycnal displacement depth of about 180 m (145 m?), in contrast to the model maxima at 121 meters. The difference in the two depths was thought to be explained by the non-isotropic internal wave energy transmission associated with the complex bathymetry. To explore this possibility, an energy ray pattern was constructed for the region. The ray trace was calculated such that the ray paths coincided with the observed phase structure at the mooring. This enabled us to trace back the rays to the assumed generation zone. The ray trace pattern (Figure 3) shows that the upward-propagating wave energy would be expected mainly in the depth range of approximately 70 to 220 meters.

Near the bank, our ray pattern is qualitatively inconsistent with that of Marsden (1986), which showed a downward ray orientations at his mooring.

Figure 12. The first-mode empirical orthogonal function in the [12 hour]-1 frequency band of the residual or non-model isopycnal displacement time series (see text). The first-order phase structure with a 180-degree phase shift at 130 m depth, is that of a dynamical second mode with (a) a mid-depth maxima at the 162 m depth and (b) a downward phase propagation below 130 m.

Appendix A1: Model Tidal Currents based on a harmonic analysis of a 17-day series => QUODDY5-tropic ~10-minute Surface Current Tides: @ **node 5179** close to the Marsden (1986) Georges Bank mooring site [42.200° N, 67.267°W].

	H (cm/s)	Phase (°G)	H (cm/s)	Phase (°G)		Maj (cm/s)	Orient (°T)	Phase (°G)	Ellipse
M ₂	30.0	182	22.3	10		37.3	-53	5	-14.4
N ₂	3.7	162	3.1	352		4.9	-52	166	- 7.8
S ₂	4.1	198	3.2	33		5.2	-34	24	- 7.8
O ₁	0.9	286	1.1	21		1.2	-25	39	1.4
K ₁	0.9	286	1.2	53		1.3	-34	71	2.1

Appendix A2: Model Tidal Currents based on a harmonic analysis of a 17-day series => QUODDY5-tropic 10-minute Surface Current Tides: @ **node 4429** close to the Brown & Irish (1993) Georges Basin mooring site [42.540°N, 67.253°W].

	H (cm/s)	Phase (°G)	H (cm/s)	Phase (°G)		Maj (cm/s)	Orient (°T)	Phase (°G)	Ellipse
M ₂	24.5	195	21.0	4		32.2	-50	10	10.7
N ₂	3.1	180	2.8	349		4.2	-48	175	10.0
S ₂	3.4	218	3.0	33		4.5	-48	36	24.0
O ₁	1.3	303	1.0	72		1.5	-57	106	2.2
K ₁	1.0	310	1.1	80		1.3	-39	101	2.2

Appendix A3: Model Tidal Currents based on a harmonic analysis of a 17-day series => QUODDY5-tropic 10-minute Surface Current Tides: @ **node 4870** [42.77°N, 67.40 °W].

	H (cm/s)	Phase (°G)	H (cm/s)	Phase (°G)		Maj (cm/s)	Orient (°T)	Phase (°G)	Ellipse
M ₂	22.1	179	33.0	10		40.0	-34	6	-11.7
N ₂	2.8	160	4.7	357		5.4	-30	173	- 7.9
S ₂	3.0	198	5.2	43		5.9	-29	37	- 5.4
O ₁	0.7	278	1.3	97		1.5	-29	97	104.9
K ₁	0.7	274	1.4	88		1.6	-26	89	21.9

References

- Brown, W.S., 1984. A comparison of Georges Bank, Gulf of Maine and New England shelf tidal dynamics, *Journal of Physical Oceanography*, Vol. 14, 145-167.
- Brown, W.S., and J.D. Irish, 1993. The annual variation of water mass structure in the Gulf of Maine, *Journal of Marine Research*, 51, 53-107.
- Brown, W.S., and J.D. Irish, 1992. The annual evolution of geostrophic flow in the Gulf of Maine, *Journal of Physical Oceanography*, Vol. 22, No. 5, 445-473.
- Chereskin, T., 1983. Generation of internal waves in Massachusetts Bay, *J. Geophys. Res.*, 88, 2649-2661.
- Garrett, C. J. R., 1972. Tidal Resonance in the Bay of Fundy and Gulf of Maine, *Nature*, vol. 238, 441-443.
- Garrett, C. J. R., and W. H. Munk, 1972. Space-time scales of internal waves, *Geophys. Fluid Dynamics*, 2, 225-264.
- La Violette, P.E., D.R. Johnson, D.A. Brooks, 1990. Sun-glitter photographs of Georges Bank and the Gulf of Maine from the space shuttle. *Oceanography*, 3, 43-49, doi:10.5670/oceanog.1990.21.
- Loder, J.W., D. Brickman, and E. Horne, 1992. Detailed structure of currents and hydrography on the northern sides of Georges Bank, *J. Geophys. Res.*, 97 (C9).
- Lynch, D.R., J.T.C. Ip, C.E. Naimie, and F.E. Werner, 1996. Comprehensive coastal circulation model with application to the Gulf of Maine. *Continental Shelf Research*, 16, 875-906.
- Marsden, R.F., 1986. The internal tide on Georges Bank, *J. Mar. Res.*, 44, 35-50.
- Moody, J.A., B. Butman, R.C. Beardsley, W.S. Brown, P. Daifuku, J.D. Irish, D.A. Mayer, H.O. Mofjeld, B. Petrie, S. Ramp, P. Smith, and W.R. Wright, 1984. Atlas of Tidal Elevation and Current Observations on the Northeast American Continental Shelf and Slope. U.S. Geological Survey Bulletin 1611, 122 pp.
- Osborne, A. R., and T. L. Burch, 1980. Internal Solitons in the Andaman Sea, *Science*, Vol. 208, Issue 4443, pp. 451-460 DOI: 10.1126/science.208.4443.451.
- Sawyer, C. 1983. Tidal phase of internal-wave generation, *J. Geophys. Res.*, 88 (C4), 2642-2648.
- Tee, K., 1979. The Structure of three-dimensional tide-generated currents, Part I. Oscillating currents, *J. Phys. Oceanogr.*, 9, 930-944.
- Trask, R.P., and M.G. Briscoe, 1983. Detection of Massachusetts Bay internal waves by the Synthetic Aperture Radar (SAR) on SEASAT, *J. Geophys. Res.*, 88, 1789-1799.
- Voorhis, A.D., 1968. Measurements of vertical motion and the partition of energy in the New England slope water, *Deep Sea Res.*, 15, 599-60.
- Wallinga, J. P., and Brown, 2004. On the Internal Tide in Massachusetts Bay
- Wright, D.G. and J.W. Loder, 1985. A depth-dependent study of the topographic rectification of tidal currents., *Geophys. Astrophys Fluid Dyn.*, 31, 169-200.